

## **HISTOPATHOLOGICAL RESCUE IN DIAGNOSIS OF CLINICALLY SIMILAR FIBROMATOUS LESIONS OF THE GINGIVA-CASE REPORTS:**

**\*<sup>1</sup>Dr. C.M.ANITHA, <sup>2</sup>Prof. Dr. S.SENTHILKUMAR M.D.S , <sup>3</sup>Prof. Dr. S. RAJASEKAR M.D.S, <sup>4</sup>Prof. Dr.V,KRISHNAN M.D.S , <sup>5</sup>Prof. Dr.R.MADHAVAN NIRMAL M.D.S, <sup>6</sup>DR.ARUMUGAM M.D.S**

<sup>\*1</sup>Post Graduate Student, <sup>2</sup>Professor and Head ,Periodontics, <sup>3</sup>Professor, , <sup>4</sup> Dean &Professor ,<sup>5</sup>Professor &Head ,Oral Pathology.

Department of Periodontics, RMDCH, Annamalai University,  
Chidambaram - 608 002, Tamil Nadu, India.

### **ABSTRACT:**

Fibromatous lesions of the gingiva are among the most common group of lesions in the oral cavity. The fibromas and inflammatory hyperplasia which accounts for majority of lesions may appear clinically and histologically similar, however with few histological differences. Both these lesions are reactive but there is a controversy to whether the fibromas are reactive or neoplastic. For conclusive differential diagnosis of these lesions Biopsy and Histopathology plays an important role. Hence we are reporting two cases of fibrous gingival lesions with differential diagnosis.